

UK A ready to use wax block bait containing brodifacoum (0.005% w/w) for use as a rodenticide by the Professional Users with Demonstrated Competence (equivalent to Trained Professionals) for the control of Norway rats (*Rattus norvegicus*), roof rats (*Rattus rattus*) and house mice (*Mus musculus*) indoors, outdoors around buildings, and in sewers, for the protection of public health, stored products and materials.

It is illegal to use this product for uses or in a manner other than that prescribed on this label.

Not for sale to the general public

Active substance: Brodifacoum 0.005% w/w
(CAS-No.: 56073-10-0)

Against mice and rats (adults and juvenile)

For use by Professional Users with Demonstrated Competence only.

This product contains a bittering agent and a dye.

Authorisation Number: GB-2015-0905

Product Type 14: Rodenticides

Read inserted leaflet and instructions before use.

First aid instructions:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K₁ administered by medical/veterinary personnel only.

In case of:

Dermal exposure: wash skin with water and then with water and soap.

Eye exposure: rinse eyes with eye-rinse liquid or water, keep eye lids open at least 10 minutes.

Oral exposure: rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Hazardous to wildlife.

Poison Information: UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

Safe Disposal of the product and its packaging:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements.


Use of gloves is recommended.

For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).



Manufactured By:
Bell
LABORATORIES, INC.
6551 N Towne Rd
Windsor WI 53598 - United States
www.belllabs.com | MADE IN USA

6468-14



NET WEIGHT: 4 kg
20g Blox

UK

Authorisation holder:
Bell Laboratories UK Limited
3 Magdalene Street
Glastonbury, England
BA6 9EW
Tel: +1 608-241-0202
Email: emea@belllabs.com

IE

Authorisation holder:
Bell Laboratories Netherlands B.V.
De Cuserstraat 93
1081 CN Amsterdam
Netherlands
+31208949140
Email: emea@belllabs.com

Batch Number, Date of Manufacture and Expiry Date are printed on the packaging.

IE A ready to use wax block bait containing brodifacoum (0.005% w/w) for use as a rodenticide by the Professionals and Trained Professionals for the control of brown rats (*Rattus norvegicus*), roof rats (*Rattus rattus*) and house mice (*Mus musculus*) indoors, outdoors around buildings, and brown rats (*Rattus norvegicus*) in sewers, for the protection of public health, stored products and materials.

Use all Biocides Safely and Sustainably

It is illegal to use this product for uses or in a manner other than that prescribed on this label.

Not for sale to the general public

Professional users must present their professional number such as herd/flock HFR/HPR number at point of sale to purchase this product for use only as specified on the label.

Active substance: Brodifacoum 0.005% w/w (CAS-No.: 56073-10-0)

Against Rats and Mice (adults and juvenile)

For Professional and Trained Professional users only

This product contains a bittering agent and a dye.

Authorisation Number: IE/BPA 71094 | Product Type 14: Rodenticides

Read inserted leaflet and instructions before use.

First aid instructions:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K₁ administered by medical/veterinary personnel only.

In case of:

Dermal exposure: wash skin with water and then with water and soap.

Eye exposure: rinse eyes with eye-rinse liquid or water, keep eye lids open at least 10 minutes.

Oral exposure: rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Bait stations must be labelled with the following information: "do not move or open", "contains a rodenticide", "product name (Solo Blox) or authorisation number (IE/BPA 71094)", "active substance (Brodifacoum)" and "in case of accident, call a poison center".

Hazardous to wildlife.

Poison Information: For information or to report a poisoning incident contact the National Poisons Information Centre, Beaumont Hospital, Dublin (01-809 2166), retain the label for reference.

Disposal of product and its packaging:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements.



100625/UK06-23|IE0125 Product Code: 933200

Insert 6199-1 must be used with this label

UK 20g Blox

Active substance: Brodifacoum 0.005% w/w (CAS-No.: 56073-10-0)

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

Instructions for use:

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.

Bait should be secured so that it cannot be dragged away from the bait station.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase (EN374).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there is resistance. Consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

Application – For Use by Trained Professionals Only

Ready-to-use bait against house mice (*Mus musculus*), Norway rats (*Rattus norvegicus*), and roof rats (*Rattus rattus*), adult and juvenile.

Instructions for use in and around buildings:

For use in and around buildings in tamper-resistant bait stations or covered bait points for the protection of goods in stock, food, health and materials.

Norway Rats and Roof Rats: High Infestation: Place 1 to 3 blocks in tamper-resistant bait station or covered and protected bait point at 5 metre intervals per placement, where rats or their signs have been observed.

Norway Rats and Roof Rats: Low Infestation: Place 1 to 3 blocks in tamper-resistant bait station or covered and protected bait point at 10 metre intervals per placement, where rats or their signs have been observed.

House Mice: High Infestation: Place 1 block in a tamper-resistant bait station or covered and protected baiting point at 2 metre intervals per placement, where mice or their signs have been observed.

House Mice: Low Infestation: Place 1 block in a tamper-resistant bait station or covered and protected baiting point at 5 metre intervals per placement, where mice or their signs have been observed.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of re-invasion.

To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas non-labile to flooding.

Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.

For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species and children.

Do not use the product in pulsed baiting treatments.

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Remove the remaining product at the end of treatment period.

Do not apply this product directly in the burrows.

Instructions for use in sewers:

For use only against Norway rats (*Rattus norvegicus*), adult and juvenile.

For the application in sewer systems, the ready-to-use bait is to be anchored or applied in bait stations or covered and protected bait points in a manner which prevents the bait from getting into contact with waste water.

Do not use this product in pulsed baiting treatments.

Up to 200 g per manhole.

Baits must be applied in a way so that they do not come into contact with water and are not washed away.

Risk mitigation measures:

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.

Do not use in areas where resistance to the active substance can be suspected.

Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.

Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

Dispose of dead rodents in accordance with local requirements. Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

First aid instructions:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K₁ administered by medical/veterinary personnel only.

In case of:

Dermal exposure: wash skin with water and then with water and soap.

Eye exposure: rinse eyes with eye-rinse liquid or water, keep eye lids open at least 10 minutes.

Oral exposure: rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Hazardous to wildlife.

Poison Information: UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

Safe Disposal of the product and its packaging:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements.

Use of gloves is recommended.

For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

IE 20g Blox

Active substance: Brodifacoum 0.005% w/w (CAS-No.: 56073-10-0)

Instructions for use

For professional and trained professional users only.

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of re-invasion.

Remove food, which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name (Solo Blox) or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.

Bait should be secured so that it cannot be dragged away from the bait station.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase (EN374).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there is resistance. Consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

Application Rates:

High infestation: up to 20g bait in baiting points every 2 metres. **Low infestation:** up to 20g bait in baiting points every 5 metres.

High infestation: up to 60g bait in baiting points every 5 metres. **Low infestation:** up to 60g bait in baiting points every 10 metres

The baiting stations should be visited at least every 2 to 3 days for mice and every 5 to 7 days for rats at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies.

Re-fill bait when necessary.

Directions for use in and around buildings – Trained Professionals:

Ready-to-use bait to be used in tamper resistant bait stations or covered and protected bait points for the protection of goods in stock, food, health and materials.

For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Do not use the product in pulsed baiting treatments.

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Remove the remaining product at the end of treatment period.

Do not apply this product directly in the burrows.

At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

Remove the remaining bait or the bait stations at the end of the treatment period.

Instructions for Use for Trained Professionals Only:

For professional and trained professional users only.

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of re-invasion.

Remove food, which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name (Solo Blox) or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.

Bait should be secured so that it cannot be dragged away from the bait station.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase (EN374).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there is resistance. Consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

Application Rates:

High infestation: up to 20g bait in baiting points every 2 metres. **Low infestation:** up to 20g bait in baiting points every 5 metres.

High infestation: up to 60g bait in baiting points every 5 metres. **Low infestation:** up to 60g bait in baiting points every 10 metres

The baiting stations should be visited at least every 2 to 3 days for mice and every 5 to 7 days for rats at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies.

Re-fill bait when necessary.

Directions for use in and around buildings – Trained Professionals:

Ready-to-use bait to be used in tamper resistant bait stations or covered and protected bait points for the protection of goods in stock, food, health and materials.

For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Do not use the product in pulsed baiting treatments.

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Remove the remaining product at the end of treatment period.

Do not apply this product directly in the burrows.

At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

Remove the remaining bait or the bait stations at the end of the treatment period.

Directions for use in and around buildings - Professionals:

For professional and trained professional users only.

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of re-invasion.

Remove food, which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name (Solo Blox) or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.

Bait should be secured so that it cannot be dragged away from the bait station.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase (EN374).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there is resistance. Consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

Application Rates:

High infestation: up to 20g bait in baiting points every 2 metres. **Low infestation:** up to 20g bait in baiting points every 5 metres.

High infestation: up to 60g bait in baiting points every 5 metres. **Low infestation:** up to 60g bait in baiting points every 10 metres

The baiting stations should be visited at least every 2 to 3 days for mice and every 5 to 7 days for rats at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies.

Re-fill bait when necessary.

Directions for use in and around buildings – Trained Professionals:

Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected bait points for the protection of goods in stock, food, health and materials.

For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Do not use the product in pulsed baiting treatments.

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Remove the remaining product at the end of treatment period.

Do not apply this product directly in the burrows.

At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

Remove the remaining bait or the bait stations at the end of the treatment period.